



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

February 13, 1845.

The MARQUIS OF NORTHAMPTON, President, in the Chair.

Ἀμύρφωτα, No. 1. "On a case of Superficial Colour presented by a Homogeneous Liquid internally colourless." By Sir John Frederick William Herschel, Bart., F.R.S., &c.

The author observed that a solution of sulphate of quinine in tartaric acid, largely diluted, although perfectly transparent and colourless when held between the eye and the light, or a white object, yet exhibits in certain aspects, and under certain incidences of the light, an extremely vivid and beautiful celestial blue colour, apparently resulting from the action of the strata which the light first penetrates on entering the liquid; and which, if not strictly superficial, at least exert their peculiar power of analysing the incident rays, and dispersing those producing the observed tint, only through a very small depth within the medium. The thinnest film of the liquid seems quite as effective in producing this superficial colour as a considerable thickness.

---

February 20, 1845.

The MARQUIS OF NORTHAMPTON, President, in the Chair.

"Additional Remarks respecting the Condensation of Gases." By Michael Faraday, Esq., F.R.S. &c.

The author, suspecting the presence of nitrogen in the nitrous oxide on which he had operated, repeated his experiments with this gas, very carefully prepared from pure nitrate of ammonia, but the results still indicated the presence of a more volatile gas mixed with another less volatile. He found that olefiant gas is readily soluble in strong alcohol, æther, oil of turpentine, and other bodies of the same kind; and that, like the former gas, it seems to be of a compound nature. His experiments confirm the prevalence of the law that the force of vapour increases in a geometrical ratio for equal increments of heat, commencing at a given amount of pressure. The more volatile a body is, the more rapidly does the force of its vapour increase by an augmentation of temperature, the increase of elasticity being directly as the volatility of the substance. By further and more accurate investigation, a general law may be established for deducing, from only a single observation of the force of any given vapour in contact with its fluid, its elasticity at any other temperature.

Postscript to the Paper by Sir John F. W. Herschel, Bart., F.R.S., read at the last meeting.

The author found that neither cinchonine nor salicine, in a state of great purity, possessed, in the smallest appreciable degree, the optical property which he has shown to belong to quinine.